

SELF-ADJOINTNESS OF THE GAFFNEY LAPLACIAN ON VECTOR BUNDLES

LASHI BANDARA AND OGNJEN MILATOVIC

ABSTRACT. We study the Gaffney Laplacian on a vector bundle equipped with a compatible metric and connection over a Riemannian manifold that is possibly geodesically incomplete. Under the hypothesis that the Cauchy boundary is polar, we demonstrate the self-adjointness of this Laplacian. Furthermore, we show that negligible boundary is a necessary and sufficient condition for the self-adjointness of this operator.

CONTENTS

1. Introduction	1
Acknowledgements	3
2. Preliminaries	3
3. General results	6
4. Polar Boundary	9
5. Negligible Boundary	11
References	13

1. INTRODUCTION

The study of the essential self-adjointness of the Laplace-Beltrami operator Δ on a geodesically complete Riemannian manifold $(\mathcal{M}, g)$ in $L^2(\mathcal{M})$ and the standard L^2 -spaces of differential forms was initiated by Gaffney in [9]. Later, also in the setting of geodesically complete Riemannian manifolds, Cordes in [7] proved the essential self-adjointness of positive integer powers of the operator Δ on $C_c^\infty(\mathcal{M})$, the space of smooth compactly supported functions on $\mathcal{M}$. In contrast to Cordes' "stationary" approach, Chernoff in [6] used a wave equation method to establish the essential self-adjointness of positive integer powers of the Laplace operator on differential forms.

In [2], the first author considered the case of a vector bundle $\mathcal{V}$ equipped with a metric h and connection ∇ that are compatible, over a geodesically complete manifold. On factorising certain "density problems" in terms of a first-order operator and applying the results of Chernoff from [6], he established the density of $C_c^\infty(\mathcal{V})$ in the Sobolev space $W^{1,2}(\mathcal{V})$ defined as the closure of

$$(1) \quad S_1(\mathcal{V}) := \{u \in C^\infty \cap L^2(\mathcal{V}) : \nabla u \in C^\infty \cap L^2(T^*\mathcal{M} \otimes \mathcal{V})\}$$

Date: May 26, 2015.

2010 Mathematics Subject Classification. 58J60, 46E35, 46E40, 47F05.

Key words and phrases. Gaffney Laplacian, Bochner Laplacian, geodesically incomplete manifold, negligible boundary, essential self-adjointness, Sobolev space, polar boundary.

under the Sobolev norm $\|\cdot\|_{W^{1,2}} = \|\cdot\| + \|\nabla\cdot\|$, where $\|\cdot\|$ stands for the L^2 -norm (see §2 for details). Additionally, the author of [2] established the density of $C_c^\infty(T^*\mathcal{M} \otimes \mathcal{V})$ in the domain of the divergence operator (the adjoint of ∇ in L^2) as well as the essential self-adjointness of the Bochner Laplacian $-\operatorname{tr} \nabla^2$ on $C_c^\infty(\mathcal{V})$, the trace with respect to the metric g of the Hessian ∇^2 (c.f. Chapter 2 in [16] by Petersen).

In the context of Riemannian manifolds $(\mathcal{M}, g)$ that are possibly geodesically incomplete, Masamune in [14] studied the essential self-adjointness of the Laplace-Beltrami operator on functions. In particular, he showed that the essential self-adjointness is equivalent to the *negligible boundary property* (see §5). In the same paper, he showed that this property is equivalent to the equality $W_0^{1,2}(\mathcal{M}) = W^{1,2}(\mathcal{M})$ (where the space $W_0^{1,2}(\mathcal{M})$ is the closure of $C_c^\infty(\mathcal{M})$ with respect to the Sobolev norm).

On a related note, it turns out that the zero capacity of the Cauchy boundary of a Riemannian manifold implies $W_0^{1,2}(\mathcal{M}) = W^{1,2}(\mathcal{M})$; see [13, 14] by Masamune and [10] by Grigor'yan and Masamune. A related study of the essential self-adjointness of the sub-Laplacian on a sub-Riemannian manifold with the pseudo-0-negligible boundary property can be found in [15] by Masamune.

In the present paper, we consider a vector bundle $\mathcal{V}$ with a compatible metric h and connection ∇ over a Riemannian manifold $(\mathcal{M}, g)$ (without boundary) that is possibly geodesically incomplete. First, by studying the effect of zero capacity of the Cauchy boundary on $(\mathcal{V}, h, \nabla)$, we show that $W_0^{1,2}(\mathcal{V}) = W^{1,2}(\mathcal{V})$ (Theorem 4.3).

We then proceed to extend the notion of negligible boundary to $(\mathcal{M}, \mathcal{V}, \nabla)$. Our fundamental object of study is the operator $\Delta_G := \nabla_c^* \overline{\nabla_2}$, where ∇_c^* is the operator adjoint of $\nabla_c := \nabla|_{C_c^\infty(\mathcal{V})}$ and $\overline{\nabla_2}$ is the closure of $\nabla_2 = \nabla|_{S_1(\mathcal{V})}$ with $S_1(\mathcal{V})$ as in (1); see §2 for details. In the context of scalar-valued functions, Δ_G appears in [10] by Grigor'yan and Masamune under the name *Gaffney Laplacian*. We remark that, Gaffney actually did not study this Laplacian, but to avoid inconsistencies with the literature, we will retain this nomenclature in our more general setting. In Theorem 5.1, we show that the self-adjointness of Δ_G is equivalent to the negligible boundary property, which is, in turn, equivalent to the equality $W_0^{1,2}(\mathcal{V}) = W^{1,2}(\mathcal{V})$. Additionally, in Theorem 5.2, which is a Bochner Laplacian analogue of Theorem 3 in [14], we establish the equivalence between essential self-adjointness of the Bochner Laplacian and negligible boundary. We emphasise that our results in particular hold for the Laplacian on vectorfields, co-vector fields, and more generally tensor fields, without requiring underlying connection to be Levi-Civita.

Our analysis rests upon an application of a version of integration by parts, which enables us to establish useful links among the first-order operators entering the definitions of Dirichlet, Neumann, and Gaffney Laplacians; see Theorem 3.3 below. Other useful links among those operators are obtained by extracting information from the negligible boundary property. Additionally, in the proof of the essential self-adjointness of the Bochner Laplacian (Theorem 5.2), we adapt the heat equation method of Masamune in [14, 15] to our setting.

ACKNOWLEDGEMENTS

The first author acknowledges the support of the Australian National University and the Australian Research Council through Discovery Project DP120103692 of McIntosh and Portal, as well as his current institution, Chalmers University of Technology/University of Gothenburg, Sweden. The authors are grateful to the anonymous referee for providing us with valuable suggestions and helping us improve the presentation of the material.

2. PRELIMINARIES

Let $\mathcal{M}$ be a smooth manifold with a smooth metric g . We emphasise that $\mathcal{M}$ is a manifold *without boundary* and that g is possibly *geodesically incomplete*, by which we mean that it may be incomplete as a metric space. We denote the induced volume measure by μ_g . We assume that $\mathcal{V}$ is a smooth vector bundle over $\mathcal{M}$, equipped with a smooth metric h .

Recall that a map $\nabla : C^\infty(\mathcal{V}) \rightarrow C^\infty(T^*\mathcal{M} \otimes \mathcal{V})$ is called a *connection* if it satisfies $\nabla_{fX}v = f\nabla_Xv$ and $\nabla_X(fv) = df \otimes v + f\nabla_Xv$ where $f \in C^\infty(\mathcal{M})$, $X \in C^\infty(T\mathcal{M})$, $v \in C^\infty(\mathcal{V})$ and df is the exterior derivative of f . A connection ∇ and metric h are *compatible* if $\nabla h \equiv 0$. We fix a connection ∇ on $\mathcal{M}$ (by which we mean on $T\mathcal{M}$) and by the same symbol, we denote a connection on $\mathcal{V}$, compatible with g and h respectively. We do not assume that ∇ on $\mathcal{M}$ is *Levi-Civita*; that is, we allow for the existence of vectorfields $X, Y \in C^\infty(T\mathcal{M})$ such that $\nabla_X Y \neq [X, Y]$ (where $[\cdot, \cdot] : C^\infty(T\mathcal{M} \times T\mathcal{M}) \rightarrow C^\infty(T\mathcal{M})$ is the *Lie bracket*).

We define the Lebesgue spaces $L^p(\mathcal{V})$ for $p \geq 1$, although we only require this theory for the case $p = 2$. First, we note some measure theoretic notions. We say that a set $A \subset \mathcal{M}$ is measurable if, whenever (U, ψ) is a chart and $U \cap A \neq \emptyset$, then the set $\psi(A \cap U)$ is Lebesgue measurable. Thus, we define the space of measurable sections $\Gamma(\mathcal{V})$, where we say that a section is measurable if its coefficients are measurable when seen through trivialisations. We remark that this notion of measurability is equivalent to measurability with respect to the induced measure μ_g . See §3 of [3] for details.

For $p \in [1, \infty)$, the set $L^p(\mathcal{M}; \mathcal{V})$ (or $L^p(\mathcal{V})$ for short) is defined as the set of sections $\xi \in \Gamma(\mathcal{V})$ such that

$$\int_{\mathcal{M}} |\xi(x)|_{h(x)}^p d\mu_g(x) < \infty.$$

We bestow $L^p(\mathcal{V})$ with the norm $\|\xi\|_p = \left(\int_{\mathcal{M}} |\xi|_h^p d\mu_g\right)^{\frac{1}{p}}$. In the case of $p = \infty$, we define $L^\infty(\mathcal{V})$ to consist of sections $\xi \in \Gamma(\mathcal{V})$ such that there exists $C > 0$ with $|\xi(x)|_{h(x)} \leq C$ for x -a.e. The norm $\|\xi\|_\infty$ is the infimum over all such constants C . Each of these spaces is a Banach space (strictly speaking, modulo sections which differ on a set of measure zero). The space $L^2(\mathcal{V})$ is a Hilbert space with inner product

$$\langle \xi, \zeta \rangle = \int_{\mathcal{M}} h_x(\xi(x), \zeta(x)) d\mu_g(x)$$

for $\xi, \zeta \in L^2(\mathcal{V})$. We assume these spaces are complex valued by identifying a real space with its complexification. The *local* L^p spaces are denoted by $L_{\text{loc}}^p(\mathcal{V})$ and they contain sections $\xi \in \Gamma(\mathcal{V})$ satisfying: $\xi \in L^p(K, \mathcal{V})$ for every open $K \subset \mathcal{M}$ with $\overline{K}$ compact.

Let

$$S_k^p(\mathcal{V}) = \{u \in C^\infty \cap L^p(\mathcal{V}) : \nabla^i u \in C^\infty \cap L^p((\otimes_{j=1}^i T^* \mathcal{M}) \otimes \mathcal{V}), i = 1, \dots, k\},$$

and define $W^{k,p}(\mathcal{V})$ as the closure of $S_k^p(\mathcal{V})$ under the norm

$$\|u\|_{W^{k,p}} = \|u\| + \sum_{i=1}^k \|\nabla^i u\|.$$

Furthermore, let $W_0^{k,p}(\mathcal{V})$ denote the closure of $C_c^\infty(\mathcal{V})$ under the same norm. We define $W_c^{k,p}(\mathcal{V})$ to be sections $\xi \in W_0^{k,p}(\mathcal{V})$ with support $\text{spt } \xi$ compact. The local Sobolev spaces $W_{\text{loc}}^{k,p}(\mathcal{V})$ consist of sections $\xi \in \Gamma(\mathcal{V})$ such that $\xi \in W^{k,p}(K, \mathcal{V})$ for every open $K \subset \mathcal{M}$ with $\overline{K}$ compact. From here on, unless otherwise stated, we will be solely concerned with the case $p = 2$. Hence, we denote $S_k^2(\mathcal{V})$ by $S_k(\mathcal{V})$.

We also note the following characterisation:

$$(W) \quad W^{1,2}(\mathcal{V}) = \{u \in L^2(\mathcal{V}) : \nabla u \in L^2(T^* \mathcal{M} \otimes \mathcal{V})\}.$$

For this, we cite Theorem 2 (i) in [14], which is a version of (W) for functions. The proof generalises with only superficial modifications to the vector bundle setting as it relies purely upon the properties of Friedrichs mollification; see the proof of Proposition 1.4 in [8].

Define $\nabla_c : C_c^\infty(\mathcal{V}) \rightarrow C_c^\infty(T^* \mathcal{M} \otimes \mathcal{V})$ and $\nabla_2 : S_1(\mathcal{V}) \rightarrow C^\infty(T^* \mathcal{M} \otimes \mathcal{V}) \cap L^2(T^* \mathcal{M} \otimes \mathcal{V})$ by $\nabla_c = \nabla$ with domain $\mathcal{D}(\nabla_c) = C_c^\infty(\mathcal{V})$ and $\nabla_2 = \nabla$ with domain $\mathcal{D}(\nabla_2) = S_1(\mathcal{V})$.

In particular, the compatibility of ∇ with h induces the following adjoint formulae:

$$(N) \quad \langle u, -\text{tr } \nabla_c v \rangle = \langle \nabla_2 u, v \rangle, \quad \text{where } u \in S_1(\mathcal{V}), v \in C_c^\infty(T^* \mathcal{M} \otimes \mathcal{V}),$$

$$(D) \quad \langle w, -\text{tr } \nabla_2 z \rangle = \langle \nabla_c w, z \rangle, \quad \text{where } w \in C_c^\infty(\mathcal{V}), z \in S_1(T^* \mathcal{M} \otimes \mathcal{V}).$$

We emphasise that the compatibility of ∇ with h is essential as the results we obtain crucially depend on these adjoint formulae.

Since $C_c^\infty(\mathcal{V})$ is dense in $L^2(\mathcal{V})$, we obtain by standard theory (i.e. Theorem III.5.28 in [11] by Kato) that the operators $\nabla_c, \nabla_2, -\text{tr } \nabla_c$ and $-\text{tr } \nabla_2$ are densely-defined and closable. Hence, we define the following operators:

$$\begin{aligned} \nabla_D &:= \overline{\nabla_c}, & \nabla_N &:= \overline{\nabla_2}, \\ \text{div}_D &:= -\nabla_c^*, & \text{div}_N &:= -\nabla_2^*, \\ \text{div}_{D,-} &:= \overline{\text{tr } \nabla_2}, & \text{div}_{N,-} &:= \overline{\text{tr } \nabla_c}. \end{aligned}$$

First, observe the following.

Proposition 2.1. *The following operator inclusions hold: $\nabla_D \subset \nabla_N$, $\operatorname{div}_N \subset \operatorname{div}_D$, $\operatorname{div}_{D,-} \subset \operatorname{div}_D$, $\operatorname{div}_{N,-} \subset \operatorname{div}_N$.*

Moreover, we obtain a characterisation of the Sobolev spaces in terms of ∇_D and ∇_N .

Proposition 2.2. *The Sobolev space $W^{1,2}(\mathcal{V}) = \mathcal{D}(\nabla_N)$ and the Sobolev space $W_0^{1,2}(\mathcal{V}) = \mathcal{D}(\nabla_D)$.*

We remark that the operators div_D and div_N can be obtained as closed, densely-defined operators even if the compatibility assumption on h and ∇ is dropped. This is a consequence of the well known fact that operators ∇_c and ∇_2 are always densely-defined and closable (c.f. Proposition 2.2 in [4]). In particular, this means that Proposition 2.2 is valid even in this more general context. However, the inclusions $\operatorname{tr} \nabla_c \subset \operatorname{div}_N$ and $\operatorname{tr} \nabla_2 \subset \operatorname{div}_D$ may no longer hold. In particular, we cannot assert that div_D and div_N are differential operators.

Dropping the compatibility requirement becomes crucial when attempting to study Sobolev spaces in the setting of low-regularity metrics, for the simple reason that the metric may not be differentiable. Some initial progress in this direction can be found in [3] for the special case of Sobolev spaces of functions under so-called “rough metrics.” These considerations are beyond the scope of this paper and we will always assume compatibility between the metric and connection unless otherwise stated.

Define the following two self-adjoint operators, which we respectively call the *Dirichlet* and *Neumann* Laplacian:

$$\Delta_D := -\operatorname{div}_D \nabla_D \text{ and } \Delta_N := -\operatorname{div}_N \nabla_N.$$

On writing the *energy* associated to our Sobolev spaces, namely,

$$E_D[u] = \int_{\mathcal{M}} |\nabla_D u|^2 d\mu_g \text{ and } E_N[v] = \int_{\mathcal{M}} |\nabla_N v|^2 d\mu_g,$$

for $u \in W_0^{1,2}(\mathcal{V})$ and $v \in W^{1,2}(\mathcal{V})$, it is immediate that $W_0^{1,2}(\mathcal{V}) = \mathcal{D}(\sqrt{\Delta_D})$ and $W^{1,2}(\mathcal{V}) = \mathcal{D}(\sqrt{\Delta_N})$. Note that when $\mathcal{M} = U \subset \mathbb{R}^n$, where U is the interior of a bounded Lipschitz domain and $\mathcal{V} = \mathcal{M} \times \mathbb{C}$, the bundle of functions, then Δ_D and Δ_N denote the classical Dirichlet and Neumann Laplacians respectively. This justifies our notation and nomenclature in this more general setting.

We also consider the composition of operators $-\operatorname{tr} \nabla_2$ and ∇_2 :

$$\Delta_s := (-\operatorname{tr} \nabla_2) \nabla_2,$$

with the induced domain $\mathcal{D}(\Delta_s) = \{u \in S_1(\mathcal{V}) : \nabla u \in S_1(T^*\mathcal{M} \otimes \mathcal{V})\}$. We call this operator the *Bochner Laplacian*. Let $\Delta_{D,s}$ and $\Delta_{N,s}$ be the restrictions of Δ_D and Δ_N to $\mathcal{D}(\Delta_D) \cap C^\infty(\mathcal{V})$ and $\mathcal{D}(\Delta_N) \cap C^\infty(\mathcal{V})$ respectively. The following operator relations easily follow from definitions and Proposition 2.1:

$$(L) \quad \Delta_{D,s} \subset \Delta_s \quad \text{and} \quad \Delta_{N,s} \subset \Delta_s.$$

Furthermore, a fundamental object of our study will be the following (not necessarily symmetric) operator:

$$\Delta_G := -\operatorname{div}_D \nabla_N.$$

As aforementioned, we call this object the Gaffney Laplacian to keep consistency with the literature.

3. GENERAL RESULTS

Let us consider an additional L^2 -differential operator $\nabla_L := (-\operatorname{div}_{N,-})^*$. Recall the containment $\operatorname{div}_{N,-} \subset \operatorname{div}_N \subset \operatorname{div}_D$ from Proposition 2.1. On taking adjoints we obtain that $\nabla_D \subset \nabla_N \subset \nabla_L$. We will show that $\nabla_L = \nabla_N$, but first, we present the following abstract lemma inspired by this endeavour.

Lemma 3.1. *Let $\mathcal{H}_1$ and $\mathcal{H}_2$ be Hilbert spaces, with inner products $\langle \cdot, \cdot \rangle_1$ and $\langle \cdot, \cdot \rangle_2$ respectively. Let $\mathcal{R}_s \subset \mathcal{H}_2$ and $\mathcal{R} \subset \mathcal{H}_2$, be dense subspaces satisfying $\mathcal{R}_s \subset \mathcal{R}$. Suppose that:*

- (i) $\mathcal{R}$ is equipped with a norm $\|\cdot\|_{\mathcal{R}}$ satisfying: there exists $c > 0$ such that $\|u\|_2 \leq c\|u\|_{\mathcal{R}}$ for all $u \in \mathcal{R}$,
- (ii) the inner product $\langle \cdot, \cdot \rangle_2$ extends continuously from $\mathcal{R}_s$ to the pairing (separably continuous bilinear map)

$$\langle \cdot, \cdot \rangle : \mathcal{R} \times \mathcal{R}' \rightarrow \mathbb{C},$$

where $\mathcal{R}'$ is the dual space of $\mathcal{R}$,

- (iii) $T : \mathcal{D}(T) \subset \mathcal{H}_1 \rightarrow \mathcal{H}_2$ and $\tilde{T} : \mathcal{H}_1 \rightarrow \mathcal{R}'$ and $\tilde{T}|_{\mathcal{D}(T)} = T$,
- (iv) $S : \mathcal{D}(S) \subset \mathcal{H}_2 \rightarrow \mathcal{H}_1$, $\mathcal{R}_s \subset \mathcal{D}(S) \subset \mathcal{R}$, and $\mathcal{D}(S)$ is dense in $(\mathcal{H}_2, \|\cdot\|_2)$ as well as $(\mathcal{R}, \|\cdot\|_{\mathcal{R}})$, and
- (v) $\langle u, Sv \rangle_1 = \langle \tilde{T}u, v \rangle$ for all $u \in \mathcal{H}_1$ and $v \in \mathcal{D}(S)$.

Then, $\mathcal{D}(S^*) = \{u \in \mathcal{H}_1 : \tilde{T}u \in \mathcal{H}_2\}$.

Proof. Let $\mathcal{D} = \{u \in \mathcal{H}_1 : \tilde{T}u \in \mathcal{H}_2\}$. Combining (iii) and (v), it is easy to see that that S and T are adjoint to each other and therefore, $\mathcal{D} \subset \mathcal{D}(S^*)$.

To prove the converse, first recall that $\mathcal{D}(S^*)$ can be characterised as the set of $u \in \mathcal{H}_1$ for which $v \mapsto \langle u, Sv \rangle_1$ is continuous. Here, continuity is measured with respect to the topology induced by $\|\cdot\|_2$ on $\mathcal{H}_2$.

Fix $u \in \mathcal{D}(S^*)$. Then, $\langle u, Sv \rangle_1 = \langle \tilde{T}u, v \rangle$ by (v) and $v \mapsto \langle \tilde{T}u, v \rangle$ is continuous in $\mathcal{H}_2$. Letting f_u be this map, continuity is equivalent to $|f_u(v)| = |\langle \tilde{T}u, v \rangle| \leq C\|v\|_2$ for some $C > 0$ whenever $v \in \mathcal{D}(S)$. Since $\mathcal{D}(S)$ is dense in $\mathcal{H}_2$ by (iv), we have that $f_u(v)$ extends to the whole of $\mathcal{H}_2$ as a bounded map. That is, $f_u \in \mathcal{H}_2'$ and hence, by the Riesz representation theorem, there exists $z \in \mathcal{H}_2$ such that $f_u(v) = \langle z, v \rangle_2$ for all $v \in \mathcal{H}_2$. Now, note $|f_u(v)| \leq C\|v\|_2 \leq cC\|v\|_{\mathcal{R}}$ where the second inequality follows from (i) and hence, $\mathcal{H}_2 \cong \mathcal{H}_2' \subset \mathcal{R}'$. That is, $z \in \mathcal{R}'$. Since $\langle \cdot, \cdot \rangle$ extends

$\langle \cdot, \cdot \rangle_2$ from $\mathcal{R}_s$ via continuity, we obtain that $\langle \tilde{T}u, v \rangle = \langle z, v \rangle_2 = \langle z, v \rangle$ for all $v \in \mathcal{D}(S)$ by invoking (ii). By the density of $\mathcal{D}(S)$ in $\mathcal{R}$ guaranteed by (iv), we can deduce that $\tilde{T}u = z$ and $\tilde{T}u \in \mathcal{H}_2$ since $z \in \mathcal{H}_2$. This proves $u \in \mathcal{D}$. $\square$

We remark that the formulation of this lemma is to provide a tool to compute the domain of an operator when there are distributional tools in hand. In application, we will see that $\mathcal{R}'$ represents the space of distributional sections, $\mathcal{R}$ an appropriate Sobolev space, and $\mathcal{R}_s$, the space of compactly supported smooth sections. We have phrased this lemma in this generality in the hope that it will be useful beyond the scope of our immediate applications in this paper.

Let $\mathcal{E}'(\mathcal{V})$ be the dual space of $C^\infty(\mathcal{V})$ and $\mathcal{D}'(\mathcal{V})$ the dual space of $C_c^\infty(\mathcal{V})$. Define $W_{\text{comp}}^{1,2}(\mathcal{V}) := W_{\text{loc}}^{1,2}(\mathcal{V}) \cap \mathcal{E}'(\mathcal{V})$. The choice of the notation “comp” in the definition is because $W_{\text{comp}}^{1,2}(\mathcal{V})$ is exactly the $W^{1,2}(\mathcal{V})$ sections with compact support, which is a consequence of the fact that $\mathcal{E}'(\mathcal{V})$ is the space of compactly supported distributional sections; see Exercise 2.3.5 in [18] by van den Ban and Crainic.

Lemma 3.2. *The $L^2(\mathcal{V})$ inner product $\langle \cdot, \cdot \rangle$ extends continuously to a pairing*

$$\langle \cdot, \cdot \rangle : W_{\text{comp}}^{1,2}(\mathcal{V}) \times W_{\text{loc}}^{-1,2}(\mathcal{V}) \rightarrow \mathbb{C}$$

from $C_c^\infty(\mathcal{V})$ by continuity. Furthermore, the equality

$$(P) \quad \langle \nabla u, v \rangle = \langle u, -\text{tr } \nabla v \rangle$$

holds if one of $\text{spt } u$ or $\text{spt } v$ is compact, and either

- (i) $u \in L_{\text{loc}}^2(\mathcal{V})$ and $v \in W_{\text{loc}}^{1,2}(T^*\mathcal{M} \otimes \mathcal{V})$, or
- (ii) $u \in W_{\text{loc}}^{1,2}(\mathcal{V})$ and $v \in L_{\text{loc}}^2(T^*\mathcal{M} \otimes \mathcal{V})$.

For the statement about the pairing in Lemma 3.2, see Lemma 9.2.9 in [18]. For the equality (P), see Lemma 8.8 in the paper [5] by Braverman, the second author and Shubin. A good reference for similar results is §7 (particularly §7.7) and Theorem 7.7 in [17] by Shubin.

Theorem 3.3. *The following operator equalities hold:*

- (i) $\nabla_L = \nabla_N$
- (ii) $\text{div}_N = \text{div}_{N,-}$
- (iii) $\text{div}_D = \text{div}_{D,-}$

Proof. Recall that $\nabla_L = (-\overline{\text{tr } \nabla_c})^* = (-\text{tr } \nabla_c)^*$. First, we use Lemma 3.2 and Lemma 3.1 and show that $\mathcal{D}(\nabla_L) = \{u \in L^2(\mathcal{V}) : \nabla u \in L^2(T^*\mathcal{M} \otimes \mathcal{V})\}$. To that end, let $\mathcal{H}_1 = L^2(\mathcal{V})$, $\mathcal{H}_2 = L^2(T^*\mathcal{M} \otimes \mathcal{V})$, $S = -\text{tr } \nabla_c$ with $\mathcal{R}_s = \mathcal{D}(S) = C_c^\infty(T^*\mathcal{M} \otimes \mathcal{V})$ and $T = \nabla_2$. Now, let $\mathcal{R} = W_{\text{comp}}^{1,2}(T^*\mathcal{M} \otimes \mathcal{V})$ with norm $\|u\|_{\mathcal{R}} = \|\nabla u\| + \|u\|$ so that $\mathcal{R}' = W_{\text{loc}}^{-1,2}(T^*\mathcal{M} \otimes \mathcal{V})$ since $W_{\text{loc}}^{-1,2}$ is the dual of $W_{\text{comp}}^{1,2}$ (see Lemma 9.2.9 in [18]). On noting that $\nabla : L^2(\mathcal{V}) \rightarrow W_{\text{loc}}^{-1,2}(T^*\mathcal{M} \otimes \mathcal{V})$, we define $\tilde{T} = \nabla$. Also, $\mathcal{D}(S) = C_c^\infty(T^*\mathcal{M} \otimes \mathcal{V}) \subset W_{\text{comp}}^{1,2}(T^*\mathcal{M} \otimes \mathcal{V}) = \mathcal{R}$ and $\mathcal{R} = W_{\text{comp}}^{1,2}(T^*\mathcal{M} \otimes \mathcal{V}) \subset L^2(T^*\mathcal{M} \otimes \mathcal{V}) = \mathcal{H}_2$. Thus, we have shown that the

hypotheses (i), (iii), (iv) of Lemma 3.1 are satisfied. Furthermore, hypothesis (ii) of Lemma 3.1 is satisfied by Lemma 3.2. Finally, the fulfilment of hypothesis (v) of Lemma 3.1 follows from (P) with $u \in L^2(\mathcal{V})$ and $v \in C_c^\infty(T^*\mathcal{M} \otimes \mathcal{V})$. Thus, we conclude that $\mathcal{D}(\nabla_L) = \{u \in L^2(\mathcal{V}) : \nabla u \in L^2(T^*\mathcal{M} \otimes \mathcal{V})\}$, and the equality $\mathcal{D}(\nabla_L) = W^{1,2}(\mathcal{V})$ follows from (W). This proves property (i).

Property (i) implies $\nabla_L^* = \nabla_N^*$, which immediately gives property (ii). For property (iii), we use the L^2 -space notations $\mathcal{H}_1 = L^2(T^*\mathcal{M} \otimes \mathcal{V})$, $\mathcal{H}_2 = L^2(\mathcal{V})$, operators $S = -\nabla_c$ and $T = \text{tr} \nabla_2$, Sobolev space $\mathcal{H} = W_{\text{comp}}^{1,2}(\mathcal{V})$, and the set $\mathcal{R}_s = \mathcal{D}(S) = C_c^\infty(\mathcal{V})$. With these notations, property (iii) follows from Lemma 3.1 and Lemma 3.2 by using the same arguments as in the proof of property (i). $\square$

The following proposition is a vector bundle analogue of Lemma 3 from [14].

Proposition 3.4. *The following equalities hold:*

$$\overline{\Delta_{D,s}} = \Delta_D \quad \text{and} \quad \overline{\Delta_{N,s}} = \Delta_N.$$

Proof. As the same kind of proof applies to both equalities, we will only prove the first one. We recall (L) and apply the heat-equation method of Masamune. Since $-\Delta_D$ is a non-positive self-adjoint operator, it generates a strongly continuous contraction semigroup $(e^{-t\Delta_D})_{t \geq 0}$ on $L^2(\mathcal{V})$. Let $u \in \mathcal{D}(\Delta_D)$ be arbitrary, and consider the family $u_t := e^{-t\Delta_D}u$. By computing u_t via the functional calculus for sectorial operators, we can easily see that $e^{-t\Delta_D} \in \mathcal{D}(\Delta_D)$ (see §D in [1]). Furthermore, since Δ_D is an elliptic operator, using elliptic regularity (see Corollary 7.1(b) and Corollary 7.4 in [17]) we have $u_t \in C^\infty(\mathcal{V})$, and, therefore, $u_t \in \mathcal{D}(\Delta_{D,s})$. Moreover, we have

$$u_t \rightarrow u \quad \text{as} \quad t \rightarrow 0^+,$$

and since $e^{-t\Delta_D}$ commutes with Δ_D (the functional calculus commutes with the operator on its domain), we have

$$\Delta_{D,s}u_t = \Delta_D u_t = e^{-t\Delta_D} \Delta_D u \rightarrow \Delta_D u, \quad \text{as } t \rightarrow 0^+,$$

where both convergence relations are understood in the L^2 -sense. This proves the equality $\overline{\Delta_{D,s}} = \Delta_D$. $\square$

The next proposition is a vector bundle analogue of Lemma 3.6 (i) and (ii) from [10].

Proposition 3.5. *The following are equivalent:*

- (i) $W_0^{1,2}(\mathcal{V}) = W^{1,2}(\mathcal{V})$,
- (ii) $\Delta_D = \Delta_N$,
- (iii) Δ_G is self-adjoint.

Proof. The equivalence of (i) and (ii) is easy. If $\Delta_D = \Delta_N$, then $\nabla_N = \nabla_D$. Hence, $-\text{div}_D = \nabla_N^*$, and, therefore, $\Delta_G = \Delta_N$. Thus, Δ_G is self-adjoint. As for the remaining implication, note that $\Delta_D \subset \Delta_G$ and $\Delta_N \subset \Delta_G$. Taking adjoints and using the self-adjointness of Δ_D , Δ_N , and Δ_G , we get the equality (i). $\square$

Remark 3.6. *Note that this proposition holds even if metric compatibility between ∇ and h is dropped, upon defining div_D and div_N abstractly as adjoints of $-\nabla_D$ and $-\nabla_N$ respectively.*

4. POLAR BOUNDARY

Let $\overline{\mathcal{M}}$ be the metric completion of $\mathcal{M}$ with respect to the Riemannian distance. We define the Cauchy boundary of $\mathcal{M}$ as

$$\partial_C \mathcal{M} := \overline{\mathcal{M}} \setminus \mathcal{M}.$$

Following [10], we now give the definition of *1-capacity* on $\mathcal{M}$. Let $\mathcal{O}$ be the collection of all open sets of $\overline{\mathcal{M}}$. For a set $\Omega \in \mathcal{O}$, we define

$$\mathcal{L}(\Omega) := \{u \in W^{1,2}(\mathcal{M}) : 0 \leq u \leq 1 \text{ and } u|_{\Omega \cap \mathcal{M}} = 1\}.$$

We define the 1-capacity of $\Omega \in \mathcal{O}$ as follows:

$$\operatorname{Cap}(\Omega) := \inf_{u \in \mathcal{L}(\Omega)} \int_{\mathcal{M}} (|u|^2 + |du|^2) d\mu_g \quad \text{if } \mathcal{L}(\Omega) \neq \emptyset.$$

Furthermore, define $\operatorname{Cap}(\Omega) = \infty$ if $\mathcal{L}(\Omega) = \emptyset$ and $\operatorname{Cap}(\emptyset) = 0$. For an arbitrary set $\Sigma \subset \mathcal{M}$, define

$$\operatorname{Cap}(\Sigma) := \inf_{\Omega \in \mathcal{O}, \Sigma \subset \Omega} \operatorname{Cap}(\Omega).$$

We call Σ *polar* if $\operatorname{Cap}(\Sigma) = 0$. If $\Sigma = \emptyset$, we set $\operatorname{Cap}(\Sigma) = 0$.

We prove that polarity of the Cauchy boundary is a sufficient condition to establish $W_0^{1,2}(\mathcal{V}) = W^{1,2}(\mathcal{V})$. But first, we prove the following approximation lemma which is noteworthy in its own right.

Lemma 4.1. *The set $L^\infty \cap W^{1,2}(\mathcal{V})$ is dense in $W^{1,2}(\mathcal{V})$.*

Proof. We use the truncation procedure of Lemma 2 in [12] by Leinfelder and Simader. By definition of $W^{1,2}(\mathcal{V})$, it is enough to show that $L^\infty(\mathcal{V}) \cap W^{1,2}(\mathcal{V})$ is dense in $S_1(\mathcal{V})$ with respect to $W^{1,2}$ -norm. By Lemma 1.16 in [8] by Eichhorn, for all $v \in C^\infty(\mathcal{V})$ we have the following diamagnetic inequality

$$(\dagger) \quad |d|v|_h| \leq |\nabla v|, \quad \mu_g\text{-a.e. } x \in M,$$

where $|\cdot|_h$ is the norm with respect to the metric h of $\mathcal{V}$. In particular, $(\dagger)$ holds for all $u \in S_1(\mathcal{V})$. Now using $(\dagger)$ and **(W)** together, we conclude that $|u|_h \in W^{1,2}(\mathcal{M})$. For simplicity we will suppress h in $|u|_h$ for the remainder of the proof.

For $R > 0$ define the following family of Lipschitz functions:

$$\psi_R(t) = \begin{cases} 1, & \text{if } t \leq R; \\ \frac{R}{t}, & \text{if } t > R. \end{cases}$$

Note that $\psi'_R(t) = 0$ if $t < R$, $0 \leq \psi_R(t) \leq 1$, $|t\psi_R(t)| \leq R$ and $|t\psi'_R(t)| \leq 1$.

We now apply Theorem A in [12] to the composition $\psi_R \circ |u|$. We remark that Theorem A in [12] was proven for the composition $f \circ w$, where $f: \mathbb{R}^k \rightarrow \mathbb{R}$ is a

Lipschitz function of class $C^1(\mathbb{R}^k \setminus \Gamma)$ with a closed countable set $\Gamma \subset \mathbb{R}^k$, and where $w: \Omega \rightarrow \mathbb{R}^k$ belongs to the Sobolev space $W^{1,2}(\Omega, \mathbb{C})$ with Ω an open set in $\mathbb{R}^m$. However, the corresponding arguments can be carried over without change to the case of functions w defined on a Riemannian manifold. Hence, we obtain

$$(2) \quad d(\psi_R \circ |u|) = \psi'_R(|u|)d|u| \quad \mu_g\text{-a.e. } x \in M.$$

As in the proof of Lemma 2 in [12] we set

$$u_R := (\psi_R \circ |u|)u.$$

Clearly, $u_R \in L^\infty(\mathcal{V}) \cap W^{1,2}(\mathcal{V})$. Using Leibniz rule and (2), we have

$$\nabla u_R = (\nabla u)\psi_R(|u|) + u \otimes (\psi'_R(|u|)d|u|),$$

in the sense of distributional sections of $\mathcal{V}$.

Let χ_G denote the characteristic function of a set G . From the properties of ψ_R it follows that

$$|\nabla u_R - \nabla u| \leq (|\nabla u| + |d|u|)|\chi_{\{|u| \geq R\}}|$$

and

$$|u_R - u| \leq |u|\chi_{\{|u| \geq R\}}.$$

Therefore, as $R \rightarrow \infty$, we have $\|u_R - u\|_{W^{1,2}} \rightarrow 0$. $\square$

Remark 4.2. We note that the diamagnetic inequality (†) as proved by Eichhorn in [8] assumes the compatibility of h and ∇ . It would be interesting to know whether this inequality still holds without this assumption.

The following is then a vector-bundle analogue of Lemma 2.2 (a) in [10].

Theorem 4.3. If $\partial_C M$ is polar, then $W_0^{1,2}(\mathcal{V}) = W^{1,2}(\mathcal{V})$.

Proof. The proof mimics that of Lemma 2.2(a) in [10]. For an arbitrary $u \in W^{1,2}(\mathcal{V})$ we construct a sequence $u_j \in W_0^{1,2}(\mathcal{V})$ such that $\|u_j - u\|_{W^{1,2}} \rightarrow 0$ as $j \rightarrow \infty$. By Lemma 4.1, we assume that $u \in L^\infty(\mathcal{V}) \cap W^{1,2}(\mathcal{V})$.

Since $\text{Cap}(\partial_C M) = 0$ there exists a sequence of open sets $\Omega_k \subset \overline{M}$, $k \geq 1$, such that $\partial_C M \subset \Omega_{k+1} \subset \Omega_k$ and $\text{Cap}(\Omega_k) \rightarrow 0$ as $k \rightarrow \infty$. For each $k \geq 1$, let $\{\varphi_j^{(k)}\}_{j \geq 1}$ be a sequence of functions with the following properties: $\varphi_j^{(k)} \in \mathcal{L}(\Omega_k)$ and $\|\varphi_j^{(k)}\|_{W^{1,2}} \rightarrow \text{Cap}(\Omega_k)$, as $j \rightarrow \infty$. Define

$$\varphi_j := \varphi_j^{(j)} \quad \text{and} \quad u_j := (1 - \varphi_j)u.$$

Fix a point $x_0 \in \mathcal{M}$ and define for $r > 0$:

$$\sigma_r(x) := \min \left\{ 1, (r^{-1}(2r - d(x, x_0)))_+ \right\},$$

where $d(\cdot, \cdot)$ is the distance with respect to the metric on $\mathcal{M}$ and f_+ is the positive part of a function f .

Clearly, $\sigma_r \in W^{1,2}(\mathcal{M})$ for all $r > 0$, $\sigma_r \rightarrow 1$, and $d\sigma_r \rightarrow 0$ as $r \rightarrow \infty$ in the μ_g -a.e. sense. We define $v_{r,j} := \sigma_r u_j$ and observe that $v_{r,j} \in W_0^{1,2}(\mathcal{V})$ and $\|v_{r,j} - u_j\|_{W^{1,2}} \rightarrow 0$ as $r \rightarrow \infty$. Thus, we may assume (without loss of generality) that $u_j \in W_0^{1,2}(\mathcal{V})$.

Since $(1 - \varphi_j) \rightarrow 1$ μ_g -a.e. as $j \rightarrow \infty$, it follows that

$$\|u_j - u\|_2 \rightarrow 0, \quad \text{as } j \rightarrow \infty.$$

Finally, using the properties $\varphi_j \rightarrow 0$ μ_g -a.e., $\nabla u \in L^2(T^*\mathcal{M} \otimes \mathcal{V})$, $u \in L^\infty(\mathcal{V})$ and $\|d\varphi_j\|_2 \rightarrow 0$, we obtain

$$\|\nabla u_j\|_2 = \|(1 - \varphi_j)\nabla u - d\varphi_j \otimes u\|_2 \rightarrow \|\nabla u\|_2,$$

as $j \rightarrow \infty$. □

The following corollary then follows directly from Proposition 3.5 and Theorem 4.3.

Corollary 4.4. *If $\partial_C M$ is polar, then Δ_G is self-adjoint.*

We remark on our results here that, much in the same way that the assumption of completeness yields the essential self-adjointness of the Bochner Laplacian (or equivalently the self-adjointness of the Gaffney Laplacian) in [2], the polarity of Cauchy boundary is very much at the level of the length structure of the manifold, and it is a sufficiently strong condition to assert the self-adjointness of the Gaffney Laplacian.

5. NEGLIGIBLE BOUNDARY

Let $(\mathcal{M}, \mathcal{V}, \nabla)$ be as in §2 and recall the set

$$S_1(\mathcal{V}) = \{u \in C^\infty \cap L^2(\mathcal{V}) : \nabla u \in C^\infty \cap L^2(T^*\mathcal{M} \otimes \mathcal{V})\}.$$

We say that $(\mathcal{M}, \mathcal{V}, \nabla)$ has *negligible boundary* if

$$(NB) \quad \langle \nabla_2 u, v \rangle = \langle u, -\text{tr } \nabla_2 v \rangle, \quad \text{for all } u \in S_1(\mathcal{V}), v \in S_1(T^*\mathcal{M} \otimes \mathcal{V}).$$

This definition is analogous to the one used by Masamune in [13, 14] in the study of the self-adjointness of the Laplacian acting on functions. The term “negligible boundary” goes back to the work of Gaffney in [9]. In this section, we will illustrate the link between this geometric condition (NB) and the equality $W_0^{1,2}(\mathcal{V}) = W^{1,2}(\mathcal{V})$.

First, we consider the relationship of (NB) to operators that we have introduced previously. The following is the link between negligible boundary and the Gaffney Laplacian.

Theorem 5.1. *The operator Δ_G is self-adjoint if and only if $(\mathcal{M}, \mathcal{V}, \nabla)$ has negligible boundary.*

Proof. If Δ_G is self-adjoint, by Proposition 3.5 we get $W_0^{1,2}(\mathcal{V}) = W^{1,2}(\mathcal{V})$. Now Proposition 2.2 implies $\nabla_D = \nabla_N$. To verify the property (NB), we first approximate

$u \in S_1(\mathcal{V})$ by a sequence $u_j \in C_c^\infty(\mathcal{V})$ in $W^{1,2}$ -norm. Next, we recall the property (D) which is a consequence of the compatibility of ∇ and $\mathbf{h}$:

$$\langle \nabla_c u_j, v \rangle = \langle u_j, -\operatorname{tr} \nabla_2 v \rangle, \quad \text{for all } v \in S_1(T^*\mathcal{M} \otimes \mathcal{V}).$$

Finally, we take the limit as $j \rightarrow \infty$ on both sides, and this shows (NB).

Now assume that $(\mathcal{M}, \mathcal{V}, \nabla)$ has negligible boundary. Our goal is to show that $-\operatorname{div}_D = \nabla_2^*$. From (NB) it follows that $-\operatorname{tr} \nabla_2 \subset \nabla_2^*$, which, after taking closures, leads to $\overline{-\operatorname{tr} \nabla_2} \subset \nabla_2^*$. By Theorem 3.3 (iii), we can rewrite the last inclusion as $-\operatorname{div}_D \subset \nabla_2^*$. Additionally, by Proposition 2.1 we have $-\operatorname{div}_N \subset -\operatorname{div}_D$, that is, $\nabla_2^* \subset -\operatorname{div}_D$. Thus, we have shown that $-\operatorname{div}_D = \nabla_2^*$. Noting that $\nabla_2^* = \nabla_N^*$, we have

$$\Delta_G = \Delta_G = -\operatorname{div}_D \nabla_N = \nabla_2^* \nabla_N = \nabla_N^* \nabla_N.$$

Now the self-adjointness of Δ_G follows by von Neumann's Theorem; see Theorem V.3.24 in [11]. $\square$

The following theorem then links (NB) to the Bochner Laplacian Δ_s .

Theorem 5.2. *The operator Δ_s is essentially self-adjoint if and only if $(\mathcal{M}, \mathcal{V}, \nabla)$ has negligible boundary.*

Proof. Assume that $(\mathcal{M}, \mathcal{V}, \nabla)$ has negligible boundary. We will first show that Δ_s is symmetric. For $w, z \in \mathcal{D}(\Delta_s)$, we have

$$\langle -\operatorname{tr} \nabla_2(\nabla_2 w), z \rangle = \langle \nabla_2 w, \nabla_2 z \rangle = \langle w, -\operatorname{tr} \nabla_2(\nabla_2 z) \rangle,$$

where in the first equality we used (NB) with $u = z$ and $v = \nabla_2 w$, and in the second equality we used (NB) with $u = w$ and $v = \nabla_2 z$. This shows that Δ_s is symmetric. We now show that $\overline{\Delta_s} = \Delta_G$. Taking closures in (L) and using Proposition 3.4 we obtain

$$(3) \quad \Delta_D \subset \overline{\Delta_s} \quad \text{and} \quad \Delta_N \subset \overline{\Delta_s}.$$

Since Δ_s is symmetric, so is $\overline{\Delta_s}$. Since Δ_D and Δ_N are self-adjoint and $\overline{\Delta_s}$ is symmetric, from (3) we get $\Delta_D = \overline{\Delta_s} = \Delta_N$. Therefore, $\overline{\Delta_s}$ is self-adjoint, that is, Δ_s is essentially self-adjoint.

Now assume that Δ_s is essentially self-adjoint. Taking closures in (L) and using Proposition 3.4 we obtain (3), which leads to $\Delta_D = \overline{\Delta_s} = \Delta_N$, that is, $W_0^{1,2}(\mathcal{V}) = W^{1,2}(\mathcal{V})$. Now by Theorem 5.1 and Proposition 3.5, it follows that $(\mathcal{M}, \mathcal{V}, \nabla)$ has negligible boundary. $\square$

To summarise, we present the following list of equivalences. We note that this easily follows from Theorem 5.1, 5.2 and Proposition 3.5.

Corollary 5.3. *The following equivalences hold:*

- (i) *the triplet $(\mathcal{M}, \mathcal{V}, \nabla)$ has negligible boundary,*
- (ii) $W_0^{1,2}(\mathcal{V}) = W^{1,2}(\mathcal{V}),$

- (iii) the operator Δ_G is self-adjoint,
- (iv) the operator Δ_s is essentially self-adjoint.

We conclude this paper with the following natural questions that have risen out of our analysis.

Question 1. *Does there exist a manifold $(\mathcal{M}, g)$ and two vector bundles $(\mathcal{V}_1, h_1, \nabla_1)$ and $(\mathcal{V}_2, h_2, \nabla_2)$ with h_i and ∇_i compatible ($i = 1, 2$), so that $W_0^{1,2}(\mathcal{V}_1) = W^{1,2}(\mathcal{V}_1)$ but $W_0^{1,2}(\mathcal{V}_2) \neq W^{1,2}(\mathcal{V}_2)$?*

Question 2. *Does there exist a manifold $(\mathcal{M}, g)$ and a vector bundle $(\mathcal{V}, h, \nabla)$ with ∇ and h compatible such that either: $W_0^{1,2}(\mathcal{M}) = W^{1,2}(\mathcal{M})$ and $W_0^{1,2}(\mathcal{V}) \neq W^{1,2}(\mathcal{V})$, or, $W_0^{1,2}(\mathcal{V}) = W^{1,2}(\mathcal{V})$ and $W_0^{1,2}(\mathcal{M}) \neq W^{1,2}(\mathcal{M})$?*

Thus, it is necessary that these questions be considered only in the case that $(\mathcal{M}, g)$ is geodesically incomplete. In this situation, we do not expect the negligible boundary property for one vector bundle to necessarily follow from another. Hence, by Corollary 5.3, we at least expect Question 1 to have an affirmative answer.

REFERENCES

- [1] David Albrecht, Xuan Duong, and Alan McIntosh, *Operator theory and harmonic analysis*, Instructional Workshop on Analysis and Geometry, Part III (Canberra, 1995), Proc. Centre Math. Appl. Austral. Nat. Univ., vol. 34, Austral. Nat. Univ., Canberra, 1996, pp. 77–136. MR 1394696 (97e:47001)
- [2] Lashi Bandara, *Density problems on vector bundles and manifolds*, Proc. Amer. Math. Soc. **142** (2014), no. 8, 2683–2695. MR 3209324
- [3] Lashi Bandara, *Rough metrics on manifolds and quadratic estimates*, ArXiv e-prints (2014).
- [4] Lashi Bandara and Alan McIntosh, *The Kato square root problem on vector bundles with generalised bounded geometry*, The Journal of Geometric Analysis (2015), 1–35.
- [5] M. Braverman, O. Milatovich, and M. Shubin, *Essential selfadjointness of Schrödinger-type operators on manifolds*, Uspekhi Mat. Nauk **57** (2002), no. 4(346), 3–58. MR 1942115 (2004g:58021)
- [6] Paul R. Chernoff, *Essential self-adjointness of powers of generators of hyperbolic equations*, J. Functional Analysis **12** (1973), 401–414. MR 0369890 (51 #6119)
- [7] H. O. Cordes, *Self-adjointness of powers of elliptic operators on non-compact manifolds*, Math. Ann. **195** (1972), 257–272. MR 0292111 (45 #1198)
- [8] Jürgen Eichhorn, *Elliptic differential operators on noncompact manifolds*, Seminar Analysis of the Karl-Weierstrass-Institute of Mathematics, 1986/87 (Berlin, 1986/87), Teubner-Texte Math., vol. 106, Teubner, Leipzig, 1988, pp. 4–169. MR 1066741 (91k:58132)
- [9] Matthew P. Gaffney, *The harmonic operator for exterior differential forms*, Proc. Nat. Acad. Sci. U. S. A. **37** (1951), 48–50. MR 0048138 (13,987b)
- [10] Alexander Grigor’yan and Jun Masamune, *Parabolicity and stochastic completeness of manifolds in terms of the Green formula*, J. Math. Pures Appl. (9) **100** (2013), no. 5, 607–632. MR 3115827
- [11] Tosio Kato, *Perturbation theory for linear operators*, Classics in Mathematics, Springer-Verlag, Berlin, 1995, Reprint of the 1980 edition. MR 1335452 (96a:47025)
- [12] Herbert Leinfelder and Christian G. Simader, *Schrödinger operators with singular magnetic vector potentials*, Math. Z. **176** (1981), no. 1, 1–19. MR 606167 (82d:35073)
- [13] Jun Masamune, *Essential self-adjointness of Laplacians on Riemannian manifolds with fractal boundary*, Comm. Partial Differential Equations **24** (1999), no. 3–4, 749–757. MR 1683058 (2000m:58035)

- [14] ———, *Analysis of the Laplacian of an incomplete manifold with almost polar boundary*, Rend. Mat. Appl. (7) **25** (2005), no. 1, 109–126. MR 2142127 (2006a:58040)
- [15] ———, *Essential self-adjointness of a sublaplacian via heat equation*, Comm. Partial Differential Equations **30** (2005), no. 10-12, 1595–1609. MR 2182306 (2006h:35035)
- [16] Peter Petersen, *Riemannian geometry*, Graduate Texts in Mathematics, vol. 171, Springer-Verlag, New York, 1998. MR 1480173 (98m:53001)
- [17] M. A. Shubin, *Pseudodifferential operators and spectral theory*, second ed., Springer-Verlag, Berlin, 2001, Translated from the 1978 Russian original by Stig I. Andersson. MR 1852334 (2002d:47073)
- [18] E.P. van den Ban and M. Crainic, *Analysis on manifolds: Lecture notes for the 2009/2010 master class*, University of Utrecht, <http://www.staff.science.uu.nl/~crain101/AS-2013/main.pdf>.

LASHI BANDARA, MATHEMATICAL SCIENCES, CHALMERS UNIVERSITY OF TECHNOLOGY AND UNIVERSITY OF GOTHENBURG, SE-412 96, GOTHENBURG, SWEDEN

URL: <http://www.math.chalmers.se/~lashitha>

E-mail address: lashi.bandara@chalmers.se

OGNJEN MILATOVIC, DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NORTH FLORIDA, JACKSONVILLE, FL 32224, USA

URL: <http://www.unf.edu/~omilatov>

E-mail address: omilatov@unf.edu